

- Bull. Planktol. Japan). 1957, 4, p. 13–15 (in Japanese with English abstract).
11. Costello, M.J., Bouchet, P., Boxshall, G., Fauchald, K., Gordon, D., Hoeksema, B.W., Poore, G.C.B., van Soest, R.W.M., Stöhr, S., Walter, T.C., Vanhoorne, B., Decock, W and Appeltans, W. Global Coordination and Standardisation in Marine Biodiversity through the World Register of Marine Species (WoRMS) and Related Databases. PLoS ONE. 2013, 8 (1), e51629. <https://doi.org/10.1371/journal.pone.0051629>.
 12. Beers, J.R. Studies on the chemical composition of the major zooplankton groups in the Sargasso Sea off Bermuda. Limnol. Oceanogr. 1966, 11 (4), p. 520–528. <https://doi.org/10.4319/lo.1966.11.4.0520>.
 13. Ikeda, T. Relationship between respiration rate and body size in marine plankton animals as a function of the temperature of habitat. Bulletin of the faculty of fisheries Hokkaido University. 1970, 21 (2), p. 91–112.
 14. The Oceanographic Society of Japan. “Zooplankton. In: Coastal environmental research manual”. Tokyo, Kouseishakouseikaku, 1986, p. 184–191. (in Japanese)
 15. Uye, S., Nagano, N., Tamaki, H. Geographical and seasonal and variations in abundance, biomass and estimated production rates of microzooplankton in the Inland Sea of Japan. J. Oceanogr. 1996, 52 (6), p. 689–703. <https://doi.org/10.1007/BF02239460>.
 16. Sato, R., Tanaka, Y., Ishimaru, T. House production by *Oikopleura dioica* (Tunicata, Appendicularia) under laboratory conditions. J. Plankton. Research. 2001, 23 (4), p. 415–423. <https://doi.org/10.1093/plankt/23.4.415>.
 17. Ikeda, T. A Growth Model for a Hyperiid Amphipod *Themisto japonica* (Bovallius) in the Japan Sea, Based on Its Intermoult Period and Moult Increment. J. Oceanogr. Soc. Japan. 1990, 46 (6), P. 261–272. <https://doi.org/10.1007/BF02123502>.
 18. Ikeda, T., Imamura, A. Abundance, vertical distribution and life cycle of a hydromedusa *Aglantha digitale* in Toyama Bay, southern Japan Sea. Bull. Plankton Soc. Japan. 1996, 43 (1), p.31–43.
 19. Iguchi, N., Ikeda, T. Production, metabolism and *P:B* ratio of *Euphausia pacifica* (Crustacea: Euphausiacea) in Toyama Bay, southern Japan Sea. Plankton Biol. Ecol. 1999, 46 (1), P. 68–74.
 20. Ukai, Y., Takahashi, K.T., Fukuchi, M., Tanimura, A. Revaluation of zooplankton wet weight data of the NORPAC net samples collected in the Indian sector of the Southern Ocean. Nankyoku Shiryo (Antarctic Record). 2014, 58 (1), p. 19–41 (in Japanese with English abstract), <https://doi.org/10.15094/00009723>.

Data Citation

1. Takahashi, K.T., Makabe, R., Odate, T. Zooplankton monitoring by a twin NORPAC net during the 58th Japanese Antarctic Research Expedition in austral summer 2016-2017. 1.00, Arctic Data archive System (ADS), Japan, 2018. <https://doi.org/10.17592/001.2018060801>.