



Zooplankton monitoring using a twin NORPAC net during the 59th Japanese Antarctic Research Expedition in austral summer 2017–2018

Kunio T. TAKAHASHI^{1,2*}, and Tsuneo ODATE^{1,2}

¹ National Institute of Polar Research, Research Organization of Information and Systems,
10–3 Midori-cho, Tachikawa, Tokyo, 190-8518, Japan

² Department of Polar Science, School of Multidisciplinary Sciences, The Graduate University for
Advanced Studies (SOKENDAI), 10–3, Midori-cho, Tachikawa, Tokyo, 190-8518, Japan

*Corresponding author. Kunio T. TAKAHASHI (takahashi.kunio@nipr.ac.jp)

(Received December 23, 2019; Accepted April 2, 2020)

Abstract: As part of the monitoring program of the Japanese Antarctic Research Expedition (JARE), zooplankton sampling using a NORPAC (NORth PACific) standard net has been routinely carried out since JARE-14 (1972/73 season) to estimate the long-term trends of abundance, species composition and their temporal/spatial variation in the uppermost 150 m of the Indian Ocean sector of the Southern Ocean. Two separate monitoring surveys were completed during JARE-59 (December 2017 to March 2018). The icebreaker *Shirase* has a fixed schedule and route, traveling down longitude 110°E each December and returning along 150°E each March. Data collected on these fixed schedule seems to be ideal for long-term monitoring work in this region. We present data from its 2017/2018 voyage. In additions, separate samplings were carried out on board the T/V *Umitaka-maru* along 110°E longitude in January 2018. This report provides the latest zooplankton data. These transect data can be used as time series and/or seasonal sets over the last 45 years.

1. Background & Summary

The need to conduct monitoring of zooplankton in the Southern Ocean has long been recognized. Since zooplankton have short life spans and faster population turnover than vertebrate predators, zooplankton communities tend to change rapidly reflecting ambient environmental conditions¹. They are closer to the base of the marine food web, and they form extremely important links between phytoplankton and higher predators. Therefore, zooplankton community structure is a useful indicator of environmental variability in the Southern Ocean^{1, 2}. Moreover, long-term monitoring of zooplankton communities provide a valuable early warning of any changes in the Southern Ocean and Antarctic ecosystem^{2, 3, 4}. Numerous intensive surveys have been conducted around Antarctica and its surrounding waters to define zooplankton composition, abundances, and

community structures. However, longer-term monitoring is difficult in the remote Southern Ocean, because research opportunities are limited in both time and space.

The Japanese Antarctic Research Expedition (JARE) has been carrying out routine zooplankton monitoring in the Indian Ocean sector of the Southern Ocean every austral summer (December–March) since 1972 (JARE-14). This monitoring program is conducted from an icebreaker, which travels along much the same cruise track at approximately the same time each year. This routine sampling is ideal for long-term monitoring of this region. Although several kinds of plankton nets have been deployed from the icebreakers, vertical hauls using a NORPAC (NORth PACific) standard net have been routinely and frequently carried out to estimate the long-term trends of abundance and species composition of surface zooplankton and its spatiotemporal variability in the upper layers of the Indian Ocean sector of the Southern Ocean. Sampling was conducted from the icebreaker *Fuji* during JARE-14 until JARE-24 (1972–1983) [5, 6](#), from the icebreaker *Shirase* during JARE-25 until JARE-49 (1983–2008) [7, 8, 9](#), from RSV *Aurora Australis* during JARE-50 (2009) [10](#), and from the new icebreaker *Shirase* during JARE-51 to JARE-58 (2009–2017) [10, 11, 12, 13, 14](#). The JARE NORPAC monitoring is the longest ongoing zooplankton study within the Antarctic regions; it has been carried for the last 45 years. In addition, zooplankton monitoring has also been carried out by the T/V *Umitaka-maru* of Tokyo University of Marine Science and Technology, since the 2013/2014 season (JARE-55) as a part of the JARE monitoring program [11, 12, 13, 14](#). This report presents the data obtained from the NORPAC standard net during JARE-59 (December 2017 to March 2018).

2. Sampling Location

From JARE-14 to JARE-28 (1986/1987), NORPAC standard net sampling sites were mainly within the western part of the Indian Ocean Sector. Thereafter, sampling stations were shifted to the east where the cruise tracks of the icebreaker *Shirase* remained along the same cruise track each season, beginning with JARE-29 (1987/1988). Regular sampling has been conducted from the icebreaker *Shirase* at five stations along longitude 110°E, ranging from 40°S to 60°S in December. Likewise, sampling is carried out at five stations along 150°E, ranging from 64°S to 45°S in March on its return voyage. The same transect covered by T/V *Umitaka-maru* also comprises six stations along 110°E, ranging from 40°S to *ca.* 65°S (ice edge). Sampling data from its traverse in January 2018 are also presented herein. Typically, routine sampling stations during the two ships' surveys are located at intervals of 5 degrees of latitude. In JARE-59, sampling station L02 (45°S, 110°E) and L08 (55°S, 150°E) of the icebreaker *Shirase* were omitted because of rough seas. The locations of sampling stations of the icebreaker *Shirase* and the T/V *Umitaka-maru* during JARE-59 are shown in [Figures 1](#) and [2](#).

3. Methods

The NORPAC standard net was established as a standard method for collecting zooplankton in international cooperative surveys at an international meeting held in Honolulu in February 1956 [15](#). A twin NORPAC standard net made of nylon bolting cloth (NGG 54, mesh size 315 μm ; NXX 13, mesh size 100 μm) was used at all sampling stations ([Fig. 3](#)). The net was hauled vertically at a speed of about 1 m/s from a depth of approximately 150 m. The maximum depth reached was estimated from the wire angle and length of wire paid out. All samples obtained were immediately preserved on board in seawater containing 5%–10% buffered formalin. The volume of water filtered by each net haul was estimated using a flow-meter (Rigo Co., Ltd., Saitama, Japan) mounted at the center of the mouth ring of the net. Sampling was conducted during the daytime at almost stations, reducing the contribution of diel vertically migrating zooplankton to the biomass.

4. Data Records

Zooplankton monitoring datasets are presented in three data sheets, species/taxa list, abundance data, and wet weight data. The fields in the dataset are:

JARE number – the JARE number of this sampling season

Ship name – the name of the ship on which the sampling was conducted

Station number – the name of the station on which the sampling was conducted

Latitude – the decimal latitude of the sampling station (negative value for South)

Longitude – the decimal longitude of the sampling station (positive value for East)

Sampling season – two-year Antarctic season based around the austral summer, e.g., “2017-2018” runs from December 2017 to March 2018

Sampling year – the sampling date year

Sampling month – the sampling date month

Sampling day – the sampling date day

Sampling time – the sampling date time (UTC)

Sampling depth – the depth of sampling tow

Mesh size – the mesh size of plankton net

Estimated volume of water filtered – the estimated volume of water filtered using a flow-meter

Abundance – the abundance of each species/taxa

Total abundance – total abundance of all zooplankton in a sample

Number of species/taxa – the number of species/taxa in a sample

Wet weight – the wet weight of each category

Total wet weight – total wet weight of all zooplankton in a sample

5. Technical Validation

5-1. Zooplankton identification

Zooplankton were identified to the lowest practical taxonomic level, generally to species or genus, using a stereo-microscope. Copepodite stages of copepod species, calyptopis stages and furcilia stages of euphausiid species were subdivided from the adults. The nauplius stages of *Rhincalanus gigas* (Copepoda: Calanoida) were distinguished from other calanoid nauplii by their large size and morphological characteristics. Zooplankton abundance was converted to individuals per cubic meter of water filtered.

The species list for this dataset was checked using the Taxon Match of the World Register of Marine Species (WoRMS: <http://www.marinespecies.org/index.php>) name validation tool. WoRMS is an open-access inventory of all marine species, being >90% complete¹⁶. The tool performs a cross check of the spelling and taxonomic status of species against the WoRMS database; it returns standard taxonomic information with valid names.

5-2. Wet weight measurement

Processing of samples was carried out according to the four-step procedure outlined below and shown in [Fig. 4](#).

Step 1: the large-sized zooplankton (more than 10 mm in size) were sorted for the whole sample in the laboratory. Zooplankton were classified into seven categories, and counting number of individuals and measured the wet weight each category using an electronic balance (Sartorius Quintix124-1SJP, readability 0.1 mg).

Step 2: after removing the larger-sized zooplankton in the step 1, all other species (<10mm in size) were counted from 1/2 to 1/32 aliquots of the whole sample, and identified to the lowest taxonomic level, generally species or genus, level, using a stereo-microscope. While sorting and counting this size fraction, the wet weight of zooplankton typically reached more than 10 mg per aliquot.

Step 3: given that the wet weight of zooplankton of 1 mm size or less were hard to sort, this fraction was offered to count the individual numbers then the wet weight was estimated by using conversion factors listed in [Table 1](#).

Step 4: the weights obtained in steps 1-3 were summed to give a total wet weight per each sample.

Zooplankton abundance per haul was converted to mg m⁻³. For a detailed description of zooplankton processing for wet-weight measurements, see Ukai *et al.* (2014)²⁵.

6. Figure




Fig. 1. Locations of the icebreaker *Shirase* sampling stations during JARE-59 in 2017/2018. Stations L02 and L08 were omitted because of rough sea conditions and these were not plotted in this figure. ■: December, ▲: March.




Fig. 2. Locations of T/V *Umitaka-maru* sampling stations during JARE-59 in January 2018.



Fig. 3. The twin NORPAC standard net. Flow-meter is attached at each mouth ring of net.




Fig. 4. Four-step procedure used to measure and to estimate the wet weight of NORPAC net samples.

7. Table

Table 1. Conversion factors between Carbon weight (CW), Dry weight (DW), Wet Weight (WW), Body length (L), Bell height (BH), Bell diameter (BD), Shell length (SL), Trunk length (TL) and Body width (BW) used in this study.

Category	Species/taxa and/or Form	Developmental stage	Conversion factors of wet weight	Conversion factors of dry weight	Conversion factors of carbon weight	
Chordata	BH/BD \geq 1		WW(μ g)=DW(μ g)/0.040	\log DW($\text{mg})=-2.333+1.268\log$ BH($\text{mm})+1.125(\log$ BH($\text{mm}))^2$	[9]	
	BH/BD<1		WW(μ g)=DW(μ g)/0.043	\log DW(μ g)=-7.67+2.75 \log BD(μ m)	[5]	
	Thecosomata, Gastropoda larvae		WW(μ g)=DW(μ g)/0.256	\log DW(μ g)=-5.10+2.46 \log SL(μ m)	[5]	
Mollusca	Cavolinidae (Cone)		WW(μ g)=(3.14*BW(μ m) ² *L(μ m))/12)*10 ⁻⁶			
	Bivalve	Larvae	WW(μ g)=DW(μ g)/0.256	\log DW(μ g)=-2.70+1.47 \log SL(μ m)	[5]	
Annelida	Polychaeta	Larvae	WW(μ g)=DW(μ g)/0.097	\log DW(μ g)=-5.68+2.10 \log L(μ m)	[5]	
	Ostracoda		WW(μ g)=DW(μ g)/0.182	\log DW(μ g)=-13.77+4.99 \log SL(μ m)	[5]	
Arthropoda	Copepoda : Calanoida	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-9.59+3.41 \log L(μ m)	[5]	
	Copepoda : Cyclopoida	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-6.05+2.10 \log L(μ m)	[5]	
	Copepoda : <i>Microssetella</i>	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-7.59+2.88 \log L(μ m)	[5]	
	Copepoda : <i>Corycaeus</i>	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-6.45+2.43 \log L(μ m)	[5]	
	Copepoda : <i>Oncaea</i>	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-5.59+2.25 \log L(μ m)	[5]	
	Copepoda : Others	Adult, Copepodite	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-9.07+3.26 \log L(μ m)	[5]	
	Copepoda : Eucalanoidae	Nauplius	WW(μ g)=DW(μ g)/0.135	\log DW(μ g)=-9.59+3.41 \log L(μ m)	[5]	
	Copepoda	Other nauplii	WW(μ g)=DW(μ g)/0.135	DW(μ g)=CW(μ g)/0.457	CW($\text{ng})=1.51*10^{-5}*L(\mu\text{m})^2.94$	[6]
	Cirripedia	Cypris	WW(μ g)=DW(μ g)/0.182	\log DW(μ g)=-13.77+4.99 \log SL(μ m)	[5]	
		Nauplius	WW(μ g)=DW(μ g)/0.182	\log DW(μ g)=-6.54+2.65 \log L(μ m)	[5]	
	Amphipoda		\log WW($\text{mg})=-1.517+2.832\log$ L($\text{mm})$			
	Euphausiacea	Calyptopis, Furcilia, Adult	WW(μ g)=DW(μ g)/0.159	DW($\text{mg})=9.954*10^{-4}*L(\text{mm})^3.156$	[10]	
		Nauplius	WW(μ g)=DW(μ g)/0.159	DW(μ g)=CW(μ g)/0.407	[5]	
	Chaetognatha		WW(μ g)=DW(μ g)/0.068	\log DW(μ g)=-0.553+2.79 \log L($\text{mm})$	[5]	
	Chordata	Doliolida, Salpida	WW(μ g)=DW(μ g)/0.050	\log DW(μ g)=-6.94+2.54 \log L(μ m)	[5]	
	Appendiculata	WW(μ g)=DW(μ g)/0.050	DW(μ g)=CW(μ g)/0.442	[5]		
Others Larvae	including eggs	WW(μ g)=(3.14*BW(μ m) ² *L(μ m))/6)*10 ⁻⁶				

[1]: Wet weight was calculated from the volume of the ellipsoid body (Specific gravity = 1). [2]: Wet weight was calculated from the volume of the cone (Specific gravity = 1). [3]: ¹⁸Ikeda (1970). [5]: ¹⁸The Oceanographic Society of Japan (1986). [6]: ^{2b}Uye et al. (1996). [7]: ²¹Sato et al. (2001). [8]: ²²Ikeda (1990). [9]: ²³Ikeda and Inamura (1996). [10]: ²⁴Iguchi et al. (1999).

Author contributions

K.T. Takahashi carried out the field sampling on board the T/V *Umitaka-maru*, and performed the processing of samples and writing of the manuscript. T. Odate directed the JARE-59 monitoring program.

Acknowledgments

The authors thank Mr. Tomohide Noguchi, Marine Works Japan Ltd., for collecting samples on board the icebreaker *Shirase*. We express our heartfelt appreciation to all members of JARE-59 for their support. We also thank the officers and crew of the icebreaker *Shirase* and the TV *Umitaka-maru*.

References

1. Reid, P.C., Colebrook, J.M., Matthews, J.B.L., Aiken, J., Continuous Plankton Recorder Team. The Continuous Plankton Recorder: concepts and history, from Plankton Indicator to undulating recorders. *Progress in Oceanography*. 2003, 58, p. 117-173.
2. Hosie, G.W., Fukuchi, M., Kawaguchi, S. Development of the Southern Ocean Continuous Plankton Recorder Survey. *Progress in Oceanography*. 2003, 58 (2-4), p. 263-283.
3. Takahashi, K., Tanimura, A., Fukuchi, M. Long-term observation of zooplankton biomass in the Indian Ocean sector of the Southern Ocean. In: *Proceedings of the International Symposium on Environmental Research in Antarctica. Memoirs of the National Institute of Polar Research. Special Issue*. 1998, 52, p. 209-219.
4. Ross, R.M., Quetin, L.B., Martinson, D.G., Iannuzzi, R.A., Stammerjohn, S.E., Smith, R.C. Palmer LTER: Patterns of distribution of five dominant zooplankton species in the epipelagic zone west of the Antarctic Peninsula, 1993-2004. *Deep-Sea Research II*. 2008, 55, p. 2086-2105.
5. Fukuchi, M., Tanimura, A. Plankton samplings on board *Fuji* in 1972–1980. *JARE data reports*. 1981, 60 (Marine biology 1), p. 1–27.
6. Watanabe, K., Nakajima, Y., Ino, Y., Sasaki, H., Fukuchi, M. Plankton samplings on board the *Fuji* in 1980–1983. *JARE data reports*. 1984, 90 (Marine biology 5), p. 1–11.
7. Takahashi, K., Tanimura, A., Fukuchi, M. Plankton sampling on board *Shirase* in 1983–1996 — NORPAC standard net samples—. *JARE data reports*. 1997, 224 (Marine biology 28), p. 1–35.
8. Sawabe, E., Takahashi, K.T., Umeda, H., Fukuchi, M. Plankton sampling on board *Shirase* in 1997–2001 —NORPAC standard net samples—. *JARE data reports*. 2005, 284 (Marine biology 32), p. 10–16.
9. Takahashi, K.T., Sawabe, E., Tsujimoto, M., Fukuchi, M. Plankton sampling on board *Shirase* in 2002–2008 —NORPAC standard net samples—. *JARE data reports*. 2008, 306 (Marine biology 38), p. 1–26.
10. Takahashi, K.T., Ojima, M., Ukai, Y., Tanimura, A. Plankton sampling from the *Aurora Australis* and

- the *Shirase* in 2009–2013 —NORPAC standard net & closing net samples—. JARE data reports. 2014, 329 (Marine biology 46), p. 1–19.
11. Takahashi, K.T., Iida, T., Ojima, M., Odate, T. Zooplankton sampling during the 55th Japanese Antarctic Research Expedition in austral summer 2013–2014. JARE data reports. 2015, 336 (Mar. Biol. 49), p. 1–15.
 12. Takahashi, K.T., Takamura T.R., Iida, T., Odate, T. Zooplankton sampling during the 56th Japanese Antarctic Research Expedition in austral summer 2014–2015. JARE data reports. 2016, 351 (Mar. Biol. 59), p. 1–15.
 13. Takahashi, K.T., Takamura T.R., Makabe, R., Odate, T. Zooplankton sampling during the 57th Japanese Antarctic Research Expedition in austral summer 2015–2016. JARE data reports. 2016, 352 (Mar. Biol. 60), p. 1–16.
 14. Takahashi, K.T., Makabe, R., Odate, T. Zooplankton monitoring using a twin NORPAC net during the 58th Japanese Antarctic Research Expedition in austral summer 2016-2017. Polar Data Journal. 2019, 3, 12-21. <https://doi.org/10.20575/00000007>
 15. Motoda, S. North Pacific standard plankton net. Nihon Plankton Kenkyu Renraku Kaihou (Inform. Bull. Planktol. Japan). 1957, 4, p. 13–15 (in Japanese with English abstract).
 16. Costello, M.J., Bouchet, P., Boxshall, G., Fauchald, K., Gordon, D., Hoeksema, B.W., Poore, G.C.B., van Soest, R.W.M., Stöhr, S., Walter, T.C., Vanhoorne, B., Decock, W and Appeltans, W. Global Coordination and Standardisation in Marine Biodiversity through the World Register of Marine Species (WoRMS) and Related Databases. PLoS ONE. 2013, 8 (1), e51629. <https://doi.org/10.1371/journal.pone.0051629>.
 17. Beers, J.R. Studies on the chemical composition of the major zooplankton groups in the Sargasso Sea off Bermuda. Limnol. Oceanogr. 1966, 11 (4), p. 520–528. <https://doi.org/10.4319/lo.1966.11.4.0520>.
 18. Ikeda, T. Relationship between respiration rate and body size in marine plankton animals as a function of the temperature of habitat. Bulletin of the faculty of fisheries Hokkaido University. 1970, 21 (2), p. 91–112.
 19. The Oceanographic Society of Japan. “Zooplankton. In: Coastal environmental research manual”. Tokyo, Kouseishakouseikaku, 1986, p. 184–191. (in Japanese)
 20. Uye, S., Nagano, N., Tamaki, H. Geographical and seasonal and variations in abundance, biomass and estimated production rates of microzooplankton in the Inland Sea of Japan. J. Oceanogr. 1996, 52 (6), p. 689–703. <https://doi.org/10.1007/BF02239460>.
 21. Sato, R., Tanaka, Y., Ishimaru, T. House production by *Oikopleura dioica* (Tunicata, Appendicularia) under laboratory conditions. J. Plankton. Research. 2001, 23 (4), p. 415–423. <https://doi.org/10.1093/plankt/23.4.415>.
 22. Ikeda, T. A Growth Model for a Hyperiid Amphipod *Themisto japonica* (Bovallius) in the Japan Sea, Based on Its Intermoult Period and Moult Increment. J. Oceanogr. Soc. Japan. 1990, 46 (6), P. 261–272. <https://doi.org/10.1007/BF02123502>.
 23. Ikeda, T., Imamura, A. Abundance, vertical distribution and life cycle of a hydromedusa *Aglantha digitale* in Toyama Bay, southern Japan Sea. Bull. Plankton Soc. Japan. 1996, 43 (1), p.31–43.

24. Iguchi, N., Ikeda, T. Production, metabolism and *P:B* ratio of *Euphausia pacifica* (Crustacea: Euphausiacea) in Toyama Bay, southern Japan Sea. *Plankton Biol. Ecol.* 1999, 46 (1), P. 68–74.
25. Ukai, Y., Takahashi, K.T., Fukuchi, M., Tanimura, A. Revaluation of zooplankton wet weight data of the NORPAC net samples collected in the Indian sector of the Southern Ocean. *Nankyoku Shiryo* (Antarctic Record). 2014, 58 (1), p. 19–41 (in Japanese with English abstract), <https://doi.org/10.15094/00009723>.

Data Citation

1. Takahashi, K.T., Odate, T. Zooplankton monitoring by a twin NORPAC net during the 59th Japanese Antarctic Research Expedition in austral summer 2017-2018. Arctic Data archive System (ADS), 1.00, NIPR, 2019, <https://doi.org/10.17592/001.2019122307>